# **FACT** SHEET

### **WHAT IS NOROVIRUS**

Norovirus is a highly contagious virus that can cause a person's digestive system to become inflamed. Norovirus is sometimes referred to as "food poisoning" or the "stomach flu." Although it can cause food poisoning, it is not caused by the influenza virus.

Learn more: ecolab.com/norovirus

Norovirus is the **LEADING CAUSE OF FOODBORNE ILLNESS OUTBREAKS** IN THE U.S.

> Source: http://www.cdc.gov/foodsafety/ pdfs/foodborne-outbreaks-annual-re port-2014-508.pdf

### SYMPTOMS OCCUR WITHIN 24 HOURS

Source: Center for Disease Control

- Severe vomiting & diarrhea Low-grade fever & chills

Nausea

Headache or muscle aches

The amount of virus particles that fit on the head of a pin are enough to infect

**PEOPLE** 

Source: Journal of Medical Virology, August, 2008



HARD TO KILL:

- Survives on surfaces and utensils for weeks
  - Can remain infectious in food at freezing temperatures and even in some cooked foods
  - Can resist many common disinfectants and sanitizers

Source: NACMCF.2016.JFP 79(5):843

## **HOW NOROVIRUS ENTERS A RESTAURANT**





**CUSTOMERS** 







**FOOD CONTAMINATED** AT ITS SOURCE

#### **FOODS TYPICALLY IMPLICATED** INCLUDE

SALADS AND SANDWICHES OR OTHER READY-TO-EAT ITEMS, OR RAW OYSTERS HARVESTED FROM SEWAGE-CONTAMINATED GROWING WATERS.

Source: Center for Disease Control



### **HOW NOROVIRUS SPREADS**

Noroviruses are found in feces or vomit of infected people and can be spread easily.





**TOUCHING** CONTAMINATED PEOPLE, OBJECTS OR SURFACES



BREATHING AIRBORNE VIRUS PARTICLES

Follow us at #foodsafetymatters (f)









**7 Learn more:** ecolab.com/nrashow



Brought to you by Ecolab, in partnership with the National Restaurant Association Restaurant.org

